



PhosphoSolutions®
Antibodies that work™

Colorado Biosciences Park
12635 East Montview Boulevard, #213
Aurora, CO 80045-7337
Tel: (888) 442-7100

Anti-GABA_A Receptor, α₃-Subunit

Catalog Number: 824-GA3N

Size: 100 μl

Product Description: Affinity purified rabbit polyclonal antibody


Applications: **WB:** 1:1000 **IHC:** 1:100

Antigen: Fusion protein from N-terminal region of the α₃-subunit of rat GABA_A receptor.

Species reactivity: The antibody has been directly tested for reactivity in Western blots with rat and mouse tissue.

Biological Significance: *Gamma*-aminobutyric acid (GABA) is the primary inhibitory neurotransmitter in the central nervous system, causing a hyperpolarization of the membrane through the opening of a Cl⁻ channel associated with the GABA_A receptor (GABA_A-R) subtype. GABA_A-Rs are important therapeutic targets for a range of sedative, anxiolytic, and hypnotic agents and are implicated in several diseases including epilepsy, anxiety, depression, and substance abuse. The GABA_A-R is a multimeric subunit complex. To date six αs, four βs and four γs, plus alternative splicing variants of some of these subunits, have been identified (Olsen and Tobin, 1990; Whiting et al., 1999; Ogris et al., 2004). Injection in oocytes or mammalian cell lines of cRNA coding for α- and β-subunits results in the expression of functional GABA_A-Rs sensitive to GABA. However, coexpression of a γ-subunit is required for benzodiazepine modulation. The various effects of the benzodiazepines in brain may also be mediated via different α-subunits of the receptor (McKernan et al., 2000; Mehta and Ticku, 1998; Ogris et al., 2004; Pörtl et al., 2003).

Anti-GABA_A-Receptor, α₃ –Subunit



Western blot of mouse brain lysates from wild type (Control) and α₃-knockout (αK/O) animals showing specific immunolabeling of the ~51k α₃-subunit of the GABA_A-R. The labeling was absent from a lysate prepared from α₃-knockout animals.

Page 1 of 2

WB = Western Blot **IF** = Immunofluorescence **IHC** = Immunohistochemistry **IP** = Immunoprecipitation

Packaging: 100 μl in 10 mM HEPES (pH 7.5), 150 mM NaCl, 100 μg per ml BSA and 50% glycerol. Adequate amount of material to conduct 10-mini Western Blots.

Storage and Stability. For long term storage –20°C is recommended. Stable at –20°C for at least 1 year.

Shipment: Domestic - Blue Ice; International – Blue Ice or Dry Ice.

Purification Method: Prepared from rabbit serum by affinity purification using a column to which the fusion protein immunogen was coupled.

Antibody Specificity: Specific for the ~51k α_3 -subunit of the GABA_A receptor in Western blots. Labeling is absent in α_3 -subunit knockout animals.

Quality Control Tests: Western blots performed on each lot.

References:

- McKernan RM, et al. (2000) Sedative but not anxiolytic properties of benzodiazepines are mediated by the GABA_A receptor α_1 -subtype. *Nature Neurosci* 3:587-592.
- Mehta AK, Ticku MK (1998) Prevalence of the GABA_A receptor assemblies containing α_1 -subunit in the rat cerebellum and cerebral cortex as determined by immunoprecipitation: Lack of modulation by chronic ethanol administration. *Mol Brain Res* 67:194-199.
- Ogris W, Pörtl A, Hauer B, Ernst M, Oberto A, Wulff P, Höger H, Wisden W, Sieghart W (2004) Affinity of various benzodiazepine site ligands in mice with a point mutation in the GABA_A receptor γ_2 -subunit. *Biochem Pharmacol* 68:1621-1629.
- Olsen RW, Tobin AJ (1990) Molecular biology of GABA_A receptors. *FASEB* 4:1469-1480.
- Pörtl A, Hauer B, Fuchs K, Tretter V, Sieghart W (2003) Subunit composition and quantitative importance of GABA_A receptor subtypes in the cerebellum of mouse and rat. *J Neurochem* 87:1444-1455.
- Whiting PJ, Bonnert TP, McKernan RM, Farrar S, Le Bourdellès B, Heavens RP, Smith DW, Hewson L, Rigby MR, Sirinathsinghji DJS, Thompson SA, Wafford KA (1999) Molecular and functional diversity of the expanding GABA_A receptor gene family. *Ann NY Acad Sci* 868:645-653.

WB = Western Blot **IF** = Immunofluorescence **IHC** = Immunohistochemistry **IP** = Immunoprecipitation

Packaging: 100 μ l in 10 mM HEPES (pH 7.5), 150 mM NaCl, 100 μ g per ml BSA and 50% glycerol. Adequate amount of material to conduct 10-mini Western Blots.

Storage and Stability. For long term storage -20°C is recommended. Stable at -20°C for at least 1 year.

Shipment: Domestic - Blue Ice; International – Blue Ice or Dry Ice.