

Anti-Phospho-Tyr¹⁴⁷² NMDA Receptor, NR2B-Subunit Antibody



PhosphoSolutions[®]
Antibodies that work™

Catalog#: p1516-1472

Size: 100 µl

www.phosphosolutions.com
orders@phosphosolutions.com
888-422-7100

Host	Applications	Species Tested	Species Reactivity*	Molecular Reference
Rabbit	WB 1:1000	M, R	B, C, Ch, H, NHP, Z	~180 kDa

Product Description: Affinity purified rabbit polyclonal antibody.

Biological Significance: The ion channels activated by glutamate that are sensitive to N-methyl-D-aspartate (NMDA) are designated NMDA receptors (NMDAR). The NMDAR plays an essential role in memory, neuronal development and it has also been implicated in several disorders of the central nervous system including Alzheimer's, epilepsy and ischemic neuronal cell death (Grosshans et al., 2002; Wenthold et al., 2003; Carroll and Zukin, 2002). The NMDA receptor is also one of the principal molecular targets for alcohol in the CNS (Lovinger et al., 1989; Alvestad et al., 2003; Snell et al., 1996). Channels with physiological characteristics are produced when the NR1 subunit is combined with one or more of the NMDAR2 (NR2 A-D) subunits (Ishii et al., 1993). Overexpression of the NR2B-subunit of the NMDA Receptor has been associated with increases in learning and memory while aged, memory impaired animals have deficiencies in NR2B expression (Clayton et al., 2002a; Clayton et al., 2002b). Recent work suggests that phosphorylation of Tyr¹⁴⁷² on NR2B may regulate the functional expression the receptor in LTP and other forms of plasticity (Nakazawa et al., 2001; Roche et al., 2001).

Antigen: Phosphopeptide corresponding to amino acid residues surrounding the phospho-Tyr¹⁴⁷² of rat NMDA NR2B.


Antibody Specificity: Specific for endogenous levels of the ~180 kDa NMDAR NR2B-subunit protein phosphorylated at Tyr¹⁴⁷². Immunolabeling is completely eliminated by treatment with λ-Ptase.

Purification Method: Prepared from pooled rabbit serum by affinity purification via sequential chromatography on phospho and non-phosphopeptide affinity columns.

Quality Control Tests: Western blots performed on each lot.

Packaging: 100 µl in 10 mM HEPES (pH 7.5), 150 mM NaCl, 100 µg BSA per ml and 50% glycerol.

Storage and Stability: Shipped on blue ice. Storage at -20°C is recommended, as aliquots may be taken without freeze/thawing due to presence of 50% glycerol. Stable for at least 1 year at -20°C.



Western blot of rat hippocampal lysate showing specific immunolabeling of the ~180 kDa NR2B subunit of the NMDAR phosphorylated at Tyr¹⁴⁷² in the first lane (-). Phosphospecificity is shown in the second lane (+) where immunolabeling is completely eliminated by lysate treatment with *lambda* phosphatase (400 units/100uL lysate for 30 min).

Application Key: WB = Western Blot IF = Immunofluorescence IHC = Immunohistochemistry IP = Immunoprecipitation

Species Reactivity Key: All-All Species A-Avian Amp-Amphibian Ar-Arabidopsis B-Bovine C-Canine Ch-Chicken D-Drosophila GP-Guinea Pig H-Human Ha-Hamster M-Mouse NHP-Non-human primate P-Pig R-Rat S-Sheep X-Xenopus Z-Zebrafish

*Species assumed based on 100% homology with sequence used as antigen

For Research Use Only

Product Specific References:

- Jang, S. S., Royston, S. E., Lee, G., Wang, S., & Chung, H. J. (2016). Seizure-Induced Regulations of Amyloid- β , STEP61, and STEP61 Substrates Involved in Hippocampal Synaptic Plasticity. *Neural plasticity*, 2016.
- Xiao, X., Levy, A. D., Rosenberg, B. J., Higley, M. J., & Koleske, A. J. (2016). Disruption of Coordinated Presynaptic and Postsynaptic Maturation Underlies the Defects in Hippocampal Synapse Stability and Plasticity in Abl2/Arg-Deficient Mice. *The Journal of Neuroscience*, 36(25), 6778-6791.
- Zamzow, D. R., Elias, V., Acosta, V. A., Escobedo, E., & Magnusson, K. R. (2016). Higher levels of phosphorylated Y1472 on GluN2B subunits in the frontal cortex of aged mice are associated with good spatial reference memory, but not cognitive flexibility. *AGE*, 38(3), 1-17.
- Jang, S. S., Royston, S. E., Xu, J., Cavaretta, J. P., Vest, M. O., Lee, K. Y., Lee, S., Jeong, H.G., Lombroso, P., & Chung, H. J. (2015). Regulation of STEP61 and tyrosine-phosphorylation of NMDA and AMPA receptors during homeostatic synaptic plasticity. *Molecular brain*, 8.
- Mao, L. M., & Wang, J. Q. (2015). Dopaminergic and cholinergic regulation of Fyn tyrosine kinase phosphorylation in the rat striatum in vivo. *Neuropharmacology*, 99, 491-499.
- Chen, W., Walwyn, W., Ennes, H. S., Kim, H., McRoberts, J. A., & Marvizón, J. C. G. (2014). BDNF released during neuropathic pain potentiates NMDA receptors in primary afferent terminals. *European Journal of Neuroscience*, 39(9), 1439-1454.
- Knox, R., Brennan-Minnella, A. M., Lu, F., Yang, D., Nakazawa, T., Yamamoto, T., Swanson, R.A., Ferriero, D.M., & Jiang, X. (2014). NR2B phosphorylation at tyrosine 1472 contributes to brain injury in a rodent model of neonatal hypoxia-ischemia. *Stroke*, 45(10), 3040-3047.
- Gladding CM, Sepers MD, Xu J, Zhang LY, Milnerwood AJ, Lombroso PJ, Raymond LA (2012) Calpain and Striatal-Enriched protein tyrosine phosphatase (STEP) activation contribute to extrasynaptic NMDA receptor localization in a Huntington's disease mouse model. *Hum Mol Genet*. Sep 1;21(17):3739-52.
- Xu J, Kurup P, Bartos JA, Patriarchi T, Hell JW, Lombroso PJ (2012) Striatal-enriched protein-tyrosine phosphatase (STEP) regulates Pyk2 kinase activity. *J Biol Chem Jun 15;287(25):20942-56*.
- Tianna R. Hicklin, Peter H. Wu, Richard A. Radcliffe, Ronald K. Freund, Susan M. Goebel-Goody, Paulo R. Correa, William R. Proctor, Paul J. Lombroso, and Michael D. Browning (2011) Alcohol inhibition of the NMDA receptor function, long-term potentiation, and fear learning requires striatal-enriched protein tyrosine phosphatase PNAS, Apr 2011; 108: 6650 - 6655.

General References:

- Alvestad RM, Grosshans DR, Coultrap SJ, Nakazawa T, Yamamoto T, Browning MD (2003) Tyrosine dephosphorylation and ethanol inhibition of N-methyl-D-aspartate receptor function. *J Biol Chem* 278:11020-11025.
- Carroll RC, Zukin RS (2002) NMDA-receptor trafficking and targeting: implications for synaptic transmission and plasticity. *Trends Neurosci* 25:571-577.
- Clayton DA, Grosshans DR, Browning MD (2002a) Aging and surface expression of hippocampal NMDA receptors. *J Biol Chem* 277:14367-14369.
- Clayton DA, Mesches MH, Alvarez E, Bickford PC, Browning MD (2002b) A hippocampal NR2B deficit can mimic age-related changes in long-term potentiation and spatial learning in the Fischer 344 rat. *J Neurosci* 22:3628-3637.
- Grosshans DR, Clayton DA, Coultrap SJ, Browning MD (2002) LTP leads to rapid surface expression of NMDA but not AMPA receptors in adult rat CA1. *Nat Neurosci* 5:27-33.
- Ishii T, Moriyoshi K, Sugihara H, Sakurada K, Kadotani H, Yokoi M, Akazawa C, Shigemoto R, Mizuno N, Masu M, Nakanishi S (1993) Molecular characterization of the family of the N-methyl- D-aspartate receptor subunits. *J Biol Chem* 268:2836-2843.
- Lovinger DM, White G, Weight FF (1989) Ethanol inhibits NMDA-activated ion current in hippocampal neurons. *Science* 243:1721-1724.
- Nakazawa T, Komai S, Tezuka T, Hisatsune C, Umemori H, Semba K, Mishina M, Manabe T, Yamamoto T (2001) Characterization of Fyn-mediated tyrosine phosphorylation sites on the NR2B subunit of the N-methyl-D-aspartate receptor. *J Biol Chem* 276:693-699.
- Roche KW, Standley S, McCallum J, Dune LC, Ehlers MD, Wenthold RJ (2001) Molecular determinants of NMDA receptor internalization. *Nat Neurosci* 4:794-802.
- Snell LD, Nunley KR, Lickteig RL, Browning MD, Tabakoff B, Hoffman PL (1996) Regional and subunit specific changes in NMDA receptor mRNA and immunoreactivity in mouse brain following chronic ethanol ingestion. *Mol Brain Res* 40:71-78.
- Wenthold RJ, Prybylowski K, Standley S, Sans N, Petralia RS (2003) Trafficking of NMDA receptors. *Annu Rev Pharmacol Toxicol* 43:335-358.